



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/427, C07D 417/06	A1	(11) International Publication Number: WO 00/27395 (43) International Publication Date: 18 May 2000 (18.05.00)
(21) International Application Number: PCT/US99/26565 (22) International Filing Date: 10 November 1999 (10.11.99) (30) Priority Data: 09/189,200 10 November 1998 (10.11.98) US (71) Applicant: ALTEON, INC. [US/US]; 170 Williams Drive, Ramsey, NJ 07446 (US). (72) Inventors: WAGLE, Dilip; Jeanne Marie Gardens, Apt. 7L, James Drive, Nanuet, NY 10954 (US). FANG, Sheng, Ding; 391 Franklin Turnpike, Mahwah, NJ (US). RHO, Taikyn; 698 Saddle River Road, Saddle Brook, NJ 07662 (US). EGAN, John, J.; 63 Ball Road, Mountain Lakes, NJ 07046 (US). ULRICH, Peter; 148 DeWolf Road, Old Tappan, NJ 07675 (US). CERAMI, Anthony; 525 East 72nd Street, New York, NY 10021 (US). (74) Agents: BLOOM, Allen et al.; Dechert Price & Rhoads, Princeton Pike Corporate Center, Building 3, Suite 210, 997 Lenox Drive, Lawrenceville, NJ 08640 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: REVERSING ADVANCED GLYCOSYLATION CROSS-LINKS USING HETEROCYCLIC-SUBSTITUTED THIAZOLIUM SALTS (57) Abstract <p>The present invention relates to compositions and methods for reversing advanced glycosylation end product-mediated cross-linking and protein aging. Accordingly, compositions are described which comprise thiazolium compounds substituted with heterocyclic groups which are capable of reversing the formation of advanced glycosylation end product cross-links. Both industrial and therapeutic applications for the invention are disclosed, as food spoilage and animal protein aging can be treated. Such compounds have particular application in the treatment of protein aging such as is responsible for the complications of aging and diabetes.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

REVERSING ADVANCED GLYCOSYLATION CROSS-LINKS USING HETEROCYCLIC-SUBSTITUTED THIAZOLIUM SALTS

5 Background of the Invention

The present invention relates generally to the aging of proteins resulting from their reaction with glucose and other reducing sugars, and more particularly to the reversing or cleavage of cross-links formed as a consequence of the formation of advanced glycosylation (glycation) end products.

10 The reaction between glucose and proteins has been known for some time. Its earliest manifestation was in the appearance of brown pigments during the cooking of food, which was identified by Maillard in 1912, who observed that glucose or other reducing sugars react with amino acids to form adducts that undergo a series of dehydrations and rearrangements to form stable brown pigments. Further studies have
15 suggested that stored and heat treated foods undergo nonenzymatic browning as a result of the reaction between glucose and the polypeptide chain, and that the proteins are resultantly cross-linked and correspondingly exhibit decreased bioavailability. As described in copending Application Serial No. 08/588,249, incorporated herein by reference, these reactions have a parallel *in vivo*, and have been found to occur with a
20 variety of other body proteins, such as lens crystallins, collagen and nerve proteins. These reactions are accelerated in the presence of elevated glucose levels, as occur in individuals with diabetes mellitus, but still occur *in vivo* at normal glucose levels. Termed advanced glycosylation (or glycation) end products (AGEs), the cross-linked products involving structural and other proteins within the body leads not only to
25 aberrant physico-chemical properties of, for example, connective tissue, but also results in the formation of new chemical structures which are recognized by specific receptors on various cell types and as a consequence of their recognition, initiate pathogenetic mechanisms leading to the complications of diabetes and aging.

Several successful therapeutic approaches have been achieved based upon
30 intervening in the accumulation of AGEs *in vivo*. One approach, exemplified in U.S. Patent 4,758,583, incorporated herein by reference, concerns the inhibition of the formation of AGEs from its precursors, by the administration of agents such as aminoguanidine and related compounds. By reacting with an early glycosylation product that results from the original reaction between the target protein and glucose, these

agents block the formation of AGEs and further formation of AGEs and cross-links in tissues is inhibited. Efficacy of this approach has been demonstrated in numerous animal models of diabetes and aging, including positive effects on macrovascular, renal, retinal, and neural pathology. These data have been reviewed by Vlassara et al., 1994,

- 5 "Biology of Diseases. Pathogenic effects of advanced glycosylation: biochemical, biologic and clinical implications for diabetes and aging," Laboratory Investigation 70:138-151; Brownlee, 1995, "The pathological implications of protein glycation," Clin. Invest. Med., 18:275-281; and Brownlee, 1995, "Advanced protein glycosylation in diabetes and aging," Ann. Rev. Med. 46:223-34.

- 10 In another pharmacological approach to controlling levels of AGEs in tissues, especially in those tissues in which AGE cross-links have already accumulated to levels which are responsible for subclinical or clinical pathology, administration of agents that reverse or break AGE cross-links has proven successful. As described in U.S. Patent No. 5,656,261 and copending U.S. Application Serial Nos. 08/588,249 and 08/848,776, all
15 of which are incorporated herein by reference in their entireties, agents and methods are disclosed which reverse (also termed cleave or break) existing AGE cross-links *in vitro* and *in vivo*. Studies demonstrate positive effects of such agents on cardiovascular complications related to aging which are accelerated in experimental diabetes (see Wolffenbuttel et al., 1998, "Breakers of Advanced Glycation End Products Restores
20 Large Artery Properties in Experimental Diabetes," Proc. Nat. Acad. Sci. U.S.A. 95:4630-4634). In these studies, rats diabetic for 9 weeks followed by 1 to 3 weeks administration of an AGE breaker compound resulted in reversal of diabetes-induced increases in large artery stiffness. Parameters that were improved included cardiac output, peripheral resistance, systemic arterial compliance, input impedance of the aorta,
25 and compliance of the carotid artery.


It is toward the identification of additional agents capable of reversing AGE cross-links that the present invention is directed.

Summary of the Invention

- 30 In accordance with the present invention, compounds and method are disclosed for reversing AGE cross-linking. AGE cross-linking caused by other reactive sugars present *in vivo* or in foodstuffs, including ribose, galactose and fructose would also be reversed by the methods and compositions of the present invention.

The agents useful in the present invention are members of the class of compounds known as thiazoliums, and in particular thiazolium compounds substituted with heterocyclic groups.

The agents comprise compounds having the following structural formula:



- 5 wherein R1 and R2 are independently selected from the group consisting of hydrogen and an alkyl group optionally substituted by a hydroxy group:
 Y is a group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{R}$ wherein R is a heterocyclic group other than alkylenedioxyaryl containing 4-10 ring members and 1-3 heteroatoms
 10 selected from the group consisting of oxygen, nitrogen and sulfur; said heterocyclic group optionally substituted by one or more substituents selected from the group consisting of alkyl, oxo, alkoxycarbonylalkyl, aryl, and aralkyl; and said one or more substituents optionally substituted by one or more alkyl or alkoxy groups;
 15 or a group of the formula $-\text{CH}_2\text{C}(=\text{O})-\text{NHR}'$ wherein R' is a heterocyclic group containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur; said heterocyclic group optionally substituted by one or more alkoxycarbonylalkyl groups;
 and X is a halide, tosylate, methanesulfonate or mesitylenesulfonate ion.
 20 The ability to reverse already-formed advanced glycosylation products carries with it significant implications in all applications where advanced glycation and concomitant molecular crosslinking is a serious detriment. In the area of food technology, for instance, the cleavage of cross-links would confer a reversal of the increased toughness resulting from the formation of AGEs during storage. In a preferred
 25 embodiment, the application of agents capable of reversal of the Maillard process has particular benefit *in vivo* as AGE cross-linking adversely affects several of the significant protein masses in the body, among them collagen, elastin, lens proteins, and the kidney

glomerular basement membrane. These proteins deteriorate both with age (hence the application of the term "protein aging") and more rapidly as a consequence of diabetes. Accordingly, the ability to reverse the cross-linking of these proteins and thus to reduce the amount of cross-links present between advanced glycosylation end products and
5 other proteins in the body carries the promise for treatment of the complications of diabetes and aging for instance, and thereby improving the quality and, perhaps, duration of animal and human life.

It is a yet further object of the present invention to provide agents which reverse the advanced glycosylation end products formed as a consequence of the aforesaid
10 advanced glycosylation reaction sequence by cleaving the α -dicarbonyl-based protein crosslinks present in the advanced glycosylation end products.


It is a still further object of the present invention to provide therapeutic methods of treating the adverse consequences of molecular or protein aging by resort to the aforesaid method and agents to achieve the reversal or cleavage of cross-links derived
15 from advanced glycosylation reactions.

It is a still further object of the present invention to provide compositions, including pharmaceutical compositions, incorporating the agents of the present invention.

Other objects and advantages will become apparent to those skilled in the art
20 from a consideration of the ensuing description.

Detailed Description of the Invention

In accordance with the present invention, agents, compositions including pharmaceutical compositions containing said agents and associated methods are
25 described which reverse already-formed advanced glycosylation end product-derived cross-links (AGE cross-links). Useful agents, for instance, comprise compounds having the structural formula:



- wherein R^1 and R^2 are independently selected from the group consisting of hydrogen and an alkyl group optionally substituted by a hydroxy group;
- Y is a group of the formula $-CH_2C(=O)R$ wherein R is a heterocyclic group other than alkylenedioxyaryl containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur; said heterocyclic group optionally substituted by one or more substituents selected from the group consisting of alkyl, oxo, alkoxycarbonylalkyl, aryl, and aralkyl groups; and said one or more substituents optionally substituted by one or more alkyl or alkoxy groups;
- or a group of the formula $-CH_2C(=O)-NHR'$ wherein R' is a heterocyclic group containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur; said heterocyclic group optionally substituted by one or more alkoxycarbonylalkyl groups;
- and X is a halide, tosylate, methanesulfonate or mesitylenesulfonate ion.

The heterocyclic groups referred to herein include 4-8 membered rings having at least one and up to 3 heteroatoms, e.g., oxygen, nitrogen, or sulfur, therein, and including various degrees of unsaturation. Representatives of such heterocyclic groups are those such as isoxazolyl, phenylisoxazolyl, furanyl, morpholino, thiomorpholino, pyrimidinyl, piperidino, homopiperidino, piperazino, methylpiperazino, hexamethyleneimino, tetrahydroquinolyl, pyridyl, methylpyridyl, imidazolyl, pyrrolidinyl, 2,6-dimethylmorpholino, furfuryl, 1,2,4-triazolyl, thiazolyl, thiophenyl, thiazolinyl, methylthiazolyl, and the like. Excluded are alkylenedioxyaryl substituents, which are described in copending application Serial No. 08/588,249, incorporated herein by reference. The heterocyclic groups of the present invention may be further substituted, for example, by an oxo group, to form, for example, a 2-oxo-tetrahydroquinolyl group,

or substituted by one or more alkyl, alkoxy, carbonylalkyl, aryl, or aralkyl groups, and such substituents may be further substituted by one or more alkyl or alkoxy groups.

Examples of Y groups of the compounds of the present include but are not limited to: 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]; 3-(2-(4-morpholinyl)-2-oxoethyl); 5 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]; 3-(2-(1-piperidinyl)-2-oxoethyl); 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinolinyl)-2-oxoethyl]; 3-(2-(1-pyrrolidinyl)-2-oxoethyl); 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]; 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]; 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(4-thiomorpholinyl)-2-oxoethyl); 3-(2-(hexahydro-1-azepinyl)-2-oxoethyl); 3-[2-(4-[2-methoxyphenyl]-1-10 piperazinyl)-2-oxoethyl]; 3-(2-(octahydro-1-azocinyl)-2-oxoethyl); 3-(2-(2-pyridinyl)-2-oxoethyl); 3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]; 3-[2-(2,6-dimethyl-1-piperidinyl)-2-oxoethyl]; 3-[2-(4-benzyl-1-piperidinyl)-2-oxoethyl]; and 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl].

The alkyl groups referred to above contain one to about eighteen carbon atoms and include, for example, methyl, ethyl, propyl, butyl, pentyl, hexyl, octyl, decyl, 15 dodecyl, and octadecyl, and the corresponding branched-chain isomers thereof. Lower alkyl groups, of one to about six carbon atoms, are preferred. The alkyl groups optionally substituted by hydroxy groups include alkyl groups as hereinbefore defined substituted with a hydroxy group at any position, such as but not limited to the following 20 examples: hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 1-hydroxypropyl, 2-hydroxypropyl, 3-hydroxypropyl, 6-hydroxyhexyl, and the like. Hydroxy-substituted lower alkyl groups as defined above are preferred. Similarly, the alkoxy groups contain from one to about eighteen carbon atoms, and include, for example, methoxy, ethoxy, propoxy, butoxy, pentoxy, hexoxy, decoxy, and octadecoxy, and the corresponding 25 branched-chain isomers thereof. Lower alkoxy groups of about one to about 6 carbons, are preferred. The alkoxy, carbonylalkyl groups encompassed by the above formula include those wherein the alkoxy portion contain from one to about eighteen carbon atoms and the alkyl portion contains from 1 to about eighteen carbon atoms. Typical alkoxy, carbonyl portions are those such as acetoxy or ethanoyloxy, propanoyloxy, 30 butanoyloxy, pentanoyloxy, hexanoyloxy, decanoyloxy, and octadecanoyloxy, and the corresponding branched chain isomers thereof. The preferred alkyl portions of these molecules have from one to about six carbon atoms.

The aryl groups encompassed by the above formula are those containing 6-10 carbon atoms, such as naphthyl, phenyl and alkyl or alkoxy substituted-phenyl, e.g., toluyl and xylyl.

For the purposes of this invention, the compounds of the present invention are
5 formed as biologically and pharmaceutically acceptable salts. Useful salt forms are the halides, particularly the bromide and chloride, tosylate, methanesulfonate, and mesitylenesulfonate salts. Other related salts can be formed using similarly non-toxic, and biologically and pharmaceutically acceptable anions.

Of the compounds encompassed herein, certain substituents are preferred. For
10 instance, the compounds wherein R_1 or R_2 are hydrogen or alkyl groups are preferred. Also highly preferred are the compounds wherein Y is a 2-oxoethyl group with a heterocyclic group of thiophenyl, thiomorpholinyl, furanyl, 2-oxo-tetrahydroquinoliny, and pyrrolidinyl.

As described in the formula above, the heterocyclic group may be represented by
15 the R group of the formula $-\text{CH}_2\text{C}(=\text{O})-\text{R}$, or it may represent the R' group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{NHR}'$. Representative, non-limiting examples of compounds of the present invention are:

- 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-thiazolium bromide
- 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide
- 20 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4-methyl-5-(2-hydroxyethyl)-thiazolium bromide
- 3-[2-[4-(2-ethoxy-2-oxoethyl)-2-thiazolyl]amino-2-oxoethyl]-4,5-dimethyl-thiazolium chloride
- 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4-methyl-5-(6-hydroxyhexyl)-thiazolium
25 bromide
- 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide
- 3-(2-(1-piperidinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
- 3-(2-(2-furanyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
- 3-(2-(2-furanyl)-2-oxoethyl)-4-(2-hydroxypentyl)thiazolium bromide
- 30 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinoliny)-2-oxoethyl]-4,5-dimethylthiazolium bromide
- 3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium bromide
- 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide

- 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide
 3-(2-(2-thienyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
 3-(2-(2-thienyl)-2-oxoethyl)-4-methyl-5-hydroxyethylthiazolium bromide
 3-(2-(4-thiomorpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
 5 3-(2-(hexahydro-1-azepinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
 3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium
 chloride
 3-(2-(octahydro-1-azocinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
 3-(2-(2-pyridinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
 10 3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride
 3-[2-(2,6-dimethyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride
 3-[2-(4-benzyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride
 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride
 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4-octylthiazolium bromide
 15 3-(2-(4-morpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
 3-[2-[4-(2-ethoxy-2-oxoethyl)-2-thiazolyl]amino-2-oxoethyl]-4,5-
 dipropylthiazolium chloride
 3-(2-(4-morpholinyl)-2-oxoethyl)-4,5-dioctadecylthiazolium bromide
 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]-4,5-dipentylthiazolium bromide
 20 3-(2-(1-piperidinyl)-2-oxoethyl)-4,5-didodecylthiazolium bromide
 3-(2-(2-furanyl)-2-oxoethyl)-5-decylthiazolium bromide
 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinolinyl)-2-oxoethyl]-4,5-dioctylthiazolium
 bromide
 3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-diethylthiazolium bromide
 25 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]-4,5-dipentylthiazolium bromide
 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]-thiazolium bromide
 3-(2-(2-thienyl)-2-oxoethyl)-thiazolium bromide
 3-(2-(2-thienyl)-2-oxoethyl)-4-methyl-5-(6-hydroxyhexyl)thiazolium bromide
 3-(2-(4-thiomorpholinyl)-2-oxoethyl)thiazolium bromide
 30 3-(2-(hexahydro-1-azepinyl)-2-oxoethyl)-4,5-dioctylthiazolium bromide
 3-(2-(octahydro-1-azocinyl)-2-oxoethyl)-4,5-didecylthiazolium bromide
 3-(2-(2-pyridinyl)-2-oxoethyl)-4,5-dioctylthiazolium bromide
 3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]-4,5-dipropylthiazolium chloride

3-[2-(2,6-dimethyl-1-piperidiny)-2-oxoethyl]-4-methylthiazolium chloride

3-[2-(4-benzyl-1-piperidiny)-2-oxoethyl]-5-methylthiazolium chloride

3-[2-(4-benzyl-1-piperaziny)-2-oxoethyl]-4-octylthiazolium chloride

5 The above compounds are capable of reversing already-formed advanced glycosylation end products on proteins. The cross-linking of proteins by formation of advanced glycosylation end products contributes to the entrapment of other proteins and results in the development *in vivo* of conditions such as reduced elasticity and wrinkling of the skin, certain kidney diseases, atherosclerosis, osteoarthritis and the like. Similarly,
10 foodstuffs including plant and animal material that undergoes nonenzymatic browning deteriorates and becomes spoiled or toughened and, consequently, inedible, unpalatable or non-nutritious. Thus, the compounds employed in accordance with this invention reduce the level of the advanced glycosylation end product-associated cross-links already present in the protein material.


15 The present methods and compositions hold the promise for reversing the aging of key proteins both in animals and plants, and concomitantly, conferring both economic and medical benefits as a result thereof. In the instance of foodstuffs, the administration of the present composition holds the promise for reversing the physico-chemical changes imparted to foodstuffs on storage, such as the increased toughness of meats that occurs
20 during aging or storage.

 The therapeutic implications of the present invention relate to the reversal of the aging process which has, as indicated earlier, been identified and exemplified in the aging of key proteins by advanced glycosylation and cross-linking. Thus, body proteins, and particularly structural body proteins, such as collagen, elastin, lens proteins, nerve
25 proteins, kidney glomerular basement membranes and other extravascular matrix components would all benefit in their longevity and operation from the practice of the present invention. The present invention thus reduces the incidence of pathologies involving the entrapment of proteins by cross-linked target proteins, such as retinopathy, cataracts, diabetic kidney disease, glomerulosclerosis, peripheral vascular disease,
30 arteriosclerosis obliterans, peripheral neuropathy, stroke, hypertension, atherosclerosis, osteoarthritis, periarticular rigidity, loss of elasticity and wrinkling of skin, stiffening of joints, glomerulonephritis, etc. Likewise, all of these conditions are in evidence and tend to occur at an accelerated rate in patients afflicted with diabetes mellitus as a

consequence of their hyperglycemia. Thus, the present therapeutic method is relevant to treatment of these and related conditions in patients either of advanced age or those suffering from one of the mentioned pathologies.

Molecular cross-linking through advanced glycosylation product formation can decrease solubility of structural proteins such as collagen in vessel walls and can also trap serum proteins, such as lipoproteins to the collagen. Also, the presence of advanced glycosylation end products and associated cross-links may result in increased permeability of the endothelium and consequently covalent trapping of extravasated plasma proteins in subendothelial matrix, as well as a reduction in susceptibility of both plasma and matrix proteins to physiologic degradation by enzymes. For these reasons, the progressive occlusion of diabetic vessels induced by chronic hyperglycemia has been hypothesized to result from excessive formation of sugar-derived and particularly, glucose-derived cross-links. Such diabetic microvascular changes and microvascular occlusion can be effectively reversed by chemical reversal or cleavage of the advanced glycosylation product cross-link formation utilizing a composition and the methods of the present invention.

Certain of the compounds disclosed herein are novel:



wherein R^1 and R^2 are independently selected from the group consisting of hydrogen and an alkyl group, optionally substituted by a hydroxy group; Y is a group of the formula $-CH_2C(=O)R$ wherein R is a heterocyclic group other than alkylenedioxyaryl containing 4-10 ring members and one heteroatom selected from the group consisting of sulfur and nitrogen, or 2-3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur; said heterocyclic group optionally substituted by one or more substituents selected from the group consisting of alkyl, oxo, alkoxycarbonylalkyl, aryl, and aralkyl

groups; and said one or more substituents optionally substituted by one or more alkyl or alkoxy groups;
or a group of the formula $-\text{CH}_2\text{C}(=\text{O})-\text{NHR}'$ wherein R' is a heterocyclic group containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur; said heterocyclic group optionally substituted by one or more alkoxycarbonylalkyl groups;
and X is a halide, tosylate, methanesulfonate or mesitylenesulfonate ion.

The heterocyclic groups referred to herein include 4-8 membered rings having at least one nitrogen or sulfur atom, or 4-8 membered rings having 2-3 heteroatoms, e.g., oxygen, nitrogen, or sulfur, therein; and including various degrees of unsaturation. Representatives of such heterocyclic groups are those such as isoxazolyl, phenylisoxazolyl, morpholino, thiomorpholino, pyrimidinyl, piperidino, homopiperidino, piperazino, methylpiperazino, hexamethyleneimino, tetrahydroquinolyl, pyridyl, methylpyridyl, imidazolyl, pyrrolidinyl, 2,6-dimethylmorpholino, furfuryl, 1,2,4-triazolyl, thiazolyl, thiophenyl, thiazolinyl, methylthiazolyl, and the like. Excluded are alkylendioxyaryl substituents, which are described in co-pending application Serial No. 08/588,249, incorporated herein by reference. The heterocyclic groups of the present invention may be further substituted, for example, by an oxo group, to form, for example, a 2-oxo-tetrahydroquinolinyl group, or substituted by one or more alkyl, alkoxycarbonylalkyl, aryl, or aralkyl groups, and such substituents may be further substituted by one or more alkyl or alkoxy groups.

The alkyl groups, alkoxy groups, alkoxycarbonylalkyl groups, aryl groups, and salts are as described hereinabove.

As described in the formula above, the heterocyclic group may be represented by the R group of the formula $-\text{CH}_2\text{C}(=\text{O})-\text{R}$, or it may represent the R' group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{NHR}'$. Representative examples of novel compounds of the present invention are:

3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-thiazolium bromide
3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide
3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4-methyl-5-(2-hydroxyethyl)-thiazolium bromide

- 3-[2-[4-(2-ethoxy-2-oxoethyl)-2-thiazolyl]amino-2-oxoethyl]-4,5-dimethylthiazolium chloride
- 3-(2-(4-morpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
- 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide
- 5 3-(2-(1-piperidiny)-2-oxoethyl)-4,5-dimethylthiazolium bromide
- 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinolinyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide
- 3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
- 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide
- 10 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide
- 3-(2-(2-thienyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
- 3-(2-(2-thienyl)-2-oxoethyl)-4-methyl-5-hydroxyethylthiazolium bromide
- 3-(2-(4-thiomorpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
- 3-(2-(hexahydro-1-azepinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium bromide
- 15 3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride
- 3-(2-(octahydro-1-azocinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
- 3-(2-(2-pyridinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide
- 3-[2-(2-methyl-1-piperidiny)-2-oxoethyl]-4,5-dimethylthiazolium chloride
- 20 3-[2-(2,6-dimethyl-1-piperidiny)-2-oxoethyl]-4,5-dimethylthiazolium chloride
- 3-[2-(4-benzyl-1-piperidiny)-2-oxoethyl]-4,5-dimethylthiazolium chloride
- 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride

As described in the Background section above, studies on other AGE cross-link cleavage agents have shown the ability to reverse cardiovascular damage in an experimental model of diabetes. As diabetes is often considered to be a model of accelerated aging, especially with respect to macrovascular complications, the agents of the present invention are suitable for therapeutic use in such complications of aging, examples of which are described above.

As will be shown in the Examples below, and as cited above, compounds capable of reversing AGE cross-links show significant promise in the treatment of various diseases and complications related to protein aging. Compounds active in an *in vitro* model of AGE cross-linking in which the compounds are evaluated for their ability to reverse the covalent cross-linking of IgG to red blood cell membranes, or cleave collagen

cross-links, has corresponding in-vivo activity in a model of protein aging. As described above, such studies demonstrate positive effects of such agents on cardiovascular complications related to aging which are accelerated in experimental diabetes (see Wolffenbuttel et al., 1998, "Breakers of Advanced Glycation End Products Restores
5 Large Artery Properties in Experimental Diabetes," Proc. Nat. Acad. Sci. U.S.A. 95:4630-4634). In these studies, rats diabetic for 9 weeks followed by 1 to 3 weeks administration of an AGE breaker compound resulted in reversal of diabetes-induced increases in large artery stiffness. Parameters that were improved included cardiac output, peripheral resistance, systemic arterial compliance, input impedance of the aorta,
10 and compliance of the carotid artery. Thus, active AGE-reversing compounds of the present invention are candidates for the treatment of the complications of protein aging.

In the instance where the compositions of the present invention are utilized for *in vivo* or therapeutic purposes, it may be noted that the compounds or agents used therein are biocompatible. Various biologically or pharmaceutically acceptable salts may be
15 used, such as halides, tosylate, etc. Pharmaceutical compositions may be prepared with a therapeutically effective quantity of the agents or compounds of the present invention and may include a pharmaceutically acceptable carrier, selected from known materials utilized for this purpose. Such compositions may be prepared in a variety of forms, depending on the method of administration. Also, various pharmaceutically acceptable
20 addition salts of the compounds of the present invention may be utilized.


A liquid form would be utilized in the instance where administration is by intravenous, intramuscular or intraperitoneal injection. When appropriate, solid dosage forms such as tablets, capsules, or liquid dosage formulations such as solutions and suspensions, etc., may be prepared for oral administration. For topical or dermal
25 application to the skin or eye, a solution, a lotion or ointment may be formulated with the agent in a suitable vehicle such as water, ethanol, propylene glycol, perhaps including a carrier to aid in penetration into the skin or eye. For example, a topical preparation could include up to about 10% of the compound of Formula I. Other suitable forms for administration to other body tissues are also contemplated.

30 In the instance where the present method has therapeutic application, the animal host intended for treatment may have administered to it a quantity of one or more of the agents, in a suitable pharmaceutical form. Administration may be accomplished by known techniques, such as oral, topical and parenteral techniques such as intradermal,

subcutaneous, intravenous or intraperitoneal injection, as well as by other conventional means. Administration of the agents may take place over an extended period of time at a dosage level of, for example, up to about 30 mg/kg.

The agent of the present invention is formulated in compositions in an amount effective to inhibit and reverse the formation of advanced glycosylation end products. This amount will, of course, vary with the particular agent being utilized and the particular dosage form, but typically is in the range of 0.01% to 1.0%, by weight, of the particular formulation.

The compounds of the present invention can be prepared generally according to the methods described in Potts et al., 1976, J. Org. Chem. 41:187, and Potts et al., 1977, J. Org. Chem. 42:1648, or as shown in the following scheme wherein R is a heterocyclic group, R¹, R², and Z are as described hereinabove, and X is a halogen atom:



In the reaction scheme above, the appropriate substituted thiazole compounds of formula I is reacted with the appropriate halo compound of formula II, to afford the desired compound of the present invention; all substituents are as hereinbefore defined.

The halo reactant may be prepared by suitable techniques known in the art. For example, for the preparation of 3-(2-thiophenyl-2-oxoethyl)-4,5-dimethyl-thiazolium bromide, the reactant 2-bromothiophene is reacted with dimethylthiazole. 2-Bromothiophene may be prepared according to the method of King et al., 1964, J. Org. Chem. 29:3459, by the bromination of 2-acetylthiophene with copper (II) bromide. Specific methods are described in the examples below.

The conditions for the reaction between the halo compound and the thiazole derivative generally involve refluxing the mixture at 110°C in an oil bath for 3-7 hours with a minimum amount of solvent such as acetonitrile, or refluxing the mixture in ethanol or acetonitrile for 3-5 hours. If the halo reactant contains chlorine, the first condition is used. For a bromo compound, the second condition is preferable.

The present invention may be better understood by reference to the following non-limiting Examples, which are provided as exemplary of the invention. The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as
5 limiting the broad scope of the invention.

Example 1

Preparation of 3-(2-(1-thienyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide

This compound was prepared by reacting 4,5-dimethylthiazole with 2-bromoacetylthiophene. 2-Bromoacetylthiophene was prepared by bromination of 2-acetylthiophene with copper (II) bromide according to the method of King et al., 1964, J.
10 Org. Chem. 29:3459. 2-Acetylthiophene (6 g, 47.6 mmol) was dissolved in chloroform (60 mL) and added to a slurry of copper (II) bromide (13.5 g, 60.44 mmol) in ethyl acetate (120 mL). The mixture was refluxed for 6 hours and then filtered while still hot through a celite pad. The filtrate cake was washed with ethyl acetate and the combined
15 filtrate was evaporated to give 2-bromoacetylthiophene (9.5 g, 97.3%). The crude product was used directly in the next reaction.

A solution of 4,5-dimethylthiazole (2.2 g, 19.4 mmol) and 2-bromoacetylthiophene (4 g, 19.4 mmol) in ethyl alcohol (20 mL) was refluxed for 3 hours. It was cooled to room temperature and t-butyl methyl ether (10 mL) was added.
20 The reaction mixture was left at room temperature overnight with stirring. The white product separated and was filtered and dried. It was crystallized from ethyl alcohol (4.42 g, 73%), m.p. 203-205°C.

Example 2

Synthesis of 3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium chloride

3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-dimethylthiazolium chloride was prepared from the reaction of N-(chloroacetyl)pyrrolidine with 4,5-dimethylthiazole. N-(chloroacetyl)pyrrolidine was prepared as follows. Pyrrolidine (63.9 g, 0.9 mol) was taken in methylene chloride (640 mL) and cooled to 0°C in a salt-ice bath. To the stirred mixture was added chloroacetyl chloride (101.8 g in 450 mL of CH₂Cl₂, 0.9 mol)
25 dropwise keeping the inside temperature below 15°C. After adding the chloroacetyl chloride, the mixture was stirred for an hour at 5°C. Sodium hydroxide solution (7 M, 190 mL) was added with vigorous stirring such that the inside temperature did not
30

exceed 20°C. The mixture was stirred for 15 minutes and the aqueous layer was separated. The organic layer was washed successively with saturated sodium bicarbonate solutions (2 x 200 mL), water (1 x 200 mL) and dried over anhydrous sodium sulfate. The solvent was removed *in vacuo* and the residue was recrystallized from hexane to give 64.5 g (48.6%) of white plate crystals, m.p. 43°C.

A mixture of N-(chloroacetyl)pyrrolidine (37.3 g, 0.25 mol), 4,5-dimethylthiazole (28.69 g, 0.25 mol) and acetonitrile (17 mL) was refluxed in an oil bath at 105-110°C for 7 hours. To the reaction mixture was added acetonitrile (230 mL) and continued to heat for 20 minutes, then t-butyl methyl ether (250 mL) was added. The reaction mixture was kept at room temperature overnight. The product was filtered and washed with a mixture of t-butyl methyl ether and acetonitrile (1:1 v/v, 100 mL), and t-butyl methyl ether (150 mL) to obtain 59.64 g (90.2% yield) of white solid. The crude product (59.64 g) was dissolved in acetonitrile (350 mL) with heating, filtered, t-butyl methyl ether (350 mL) added, and the solution was allowed to cool at room temperature for 3 hours. The product was filtered and washed with a mixture of t-butyl methyl ether and acetonitrile (1:1 v/v, 300 mL), and t-butyl methyl ether (300 mL) to yield 56.95 g (86.1% yield) of white crystals, m.p. 196-198°C.

Example 3

Using the procedures described above in Examples 1 and 2, the following compounds were prepared using the corresponding reactants. The melting points are indicated.

3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-thiazolium bromide	213-214°C
3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide	245-246°C
	(dec.)
3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4-methyl-5-(2-hydroxyethyl)-thiazolium bromide	209-210°C
	(dec.)
3-[2-[4-(2-ethoxy-2-oxoethyl)-2-thiazolyl]amino-2-oxoethyl]-4,5-dimethylthiazolium chloride	120-121°C
3-(2-(4-morpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide	197-198°C
	(dec.)
3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide	215-216°C
3-(2-(1-piperidinyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide	240-242°C
3-(2-(2-furanyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide	195-196°C
	(dec.)
3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinolinyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide	274-275°C
	(dec.)

3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium chloride	189-198°C
3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide	171-172°C
3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium bromide	150-152°C
3-(2-(2-thienyl)-2-oxoethyl)-4,5-dimethyl-thiazolium bromide	203-205°C
3-(2-(2-thienyl)-2-oxoethyl)-4-methyl-5-hydroxyethylthiazolium bromide	175-176°C
3-(2-(4-thiomorpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium chloride	232-233°C
	(dec.)
3-(2-(hexahydro-1-azepinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium chloride	218-219°C
	(dec.)
3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride	215-216°C
	(dec.)
3-(2-(octahydro-1-azocinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium chloride	210-212°C
	(dec.)
3-(2-(2-pyridinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium bromide	178-179°C
3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride	220-221°C
	(dec.)
3-[2-(2,6-dimethyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride	248-249°C
	(dec.)
3-[2-(4-benzyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride	170-171°C
3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium chloride	180-182°C

Example 4

A typical pharmaceutical dosage form is prepared as follows.

	<u>mg / tablet</u>
Compound of the present invention	50
Starch	50
Mannitol	75
Magnesium stearate	2
Stearic acid	5

- 5 The compound, a portion of the starch and the lactose are combined and wet granulated with starch paste. The wet granulation is placed on trays and allowed to dry overnight at a temperature of 45°C. The dried granulation is comminuted in a comminutor to a particle size of approximately 20 mesh. Magnesium stearate, stearic acid and the balance of the starch are added and the entire mix blended prior to
- 10 compression on a suitable tablet press. The tablets are compressed at a weight of 232 mg. using a 11/32" punch with a hardness of 4 kg. These tablets will disintegrate within a half hour according to the method described in USP XVI.

Example 5

<u>Lotion</u>	<u>mg/g</u>
Compound of Formula I	1.0
Ethyl alcohol	400.0

- 18 -

Polyethylene glycol 400	300.0
Hydroxypropyl cellulose	5.0
Propylene glycol	to make 1.0 g

Example 6 - Reversing Age Cross-Links

The activity of the compounds of the present invention to reverse AGE cross-links was evaluated in an in-vitro model, as follows. Blood was collected from
5 streptozotocin-diabetic rats (6 to 8 wk diabetic) in heparinized tubes, centrifuged at 2500 X g for 15 min. at 40°C, and the plasma and buffy-coat layers aspirated off. The RBC were washed with PBS (ca. 5 ml PBS per ml blood) three times.

Test compounds were dissolved in PBS and the pH adjusted to 7.0, if necessary. Two hundred microliters of washed RBC were added to 2 mL of test compound solution,
10 and the mixture was incubated overnight at 37°C. For a control, 200 µl of RBC was incubated in 2 mL of PBS.

After the overnight incubation, the reaction mixtures were centrifuged and the RBC pellets were washed three times with PBS, followed by diluting 1:30 in PBS. Immunoglobulin G (IgG) bound to the surface of the washed RBCs was then determined.

15 Assay for RBC-IgG: The assay is performed in a Multiscreen-HA, 0.45 µm cellulose mixed esters membrane-sealed 96 well plate (Millipore MAHAS45). The membranes are first wet by filling the wells with 100 uL PBS, and the wells emptied by applying a vacuum to the Millititer vacuum manifold. Three hundred uL of Superblock (pre-warmed to 37°C) is then added to each well and incubated at 37°C for one hour.
20 The Superblock is then removed from the wells by application of vacuum, then the wells are washed once with 300 ul of PBS-Tween (0.05%) and three times with PBS. The vacuum is then turned off.

One hundred µl of PBS is then added to each well. Each RBC sample is gently vortexed, and 50 uL pipetted into the wells, in sextuplicate. The wells labeled A1-A3 are
25 left for a reagent blank and wells A4-A6 reserved for an antibody blank. Vacuum is then applied to remove the buffer and the RBCs are washed once with PBS. Alkaline phosphatase-labeled rabbit-anti-rat IgG is diluted 1 to 25000 in PBS, and 50 µl is added to triplicate wells of each sample. PBS alone is added to the other three wells of each sample. These will serve as sample blanks to adjust for any endogenous alkaline
30 phosphatase activity. The samples are allowed to stand at room temperature for two

hours. The solution is then removed by application of vacuum and the red blood cells are washed with PBS-Tween twice, PBS twice and with TBS twice. The bottom of the plate is rinsed with distilled water and blotted dry with paper towels. To each well is then added 100 μ l of p-nitrophenyl phosphate substrate (1 mg/ml in diethanolamine buffer pH 9.5), and the color is allowed to develop for two hours at 37°C. The solutions in each well are then transferred to a 96 well microtiter plate inside the vacuum manifold. One hundred μ l of PBS to each well and vacuum applied again to transfer any remaining solution into the microtiter plate. The OD at 410nm is then read in a Dynatech Plate reader (Sample filter 1 and Ref. filter 4).

10 For data analysis, breaking activity (percent reversal) is expressed as the percent decrease in O.D. caused by incubation of RBC with the test compound compared to RBC incubated in PBS alone. Using the above assay, the data was generated on compounds of the present invention.

Example 7 - Reversing Collagen Cross Links

15 In a further evaluation of the ability of compounds of the present invention to reverse AGE cross-links between a circulating protein and collagen, AGE-BSA was cross-linked to rat tail tendon collagen (type 1)-coated 96 well microtiter plates, and reversal of cross-linking was determined.

AGE-BSA preparation. AGE-BSA is prepared by incubation of a solution of BSA (400 mg/ml) in 0.4 M sodium phosphate buffer pH 7.4 with an equal volume of 400 mM glucose solution in the same buffer at 37°C for 12 weeks. The pH of the incubation mixture is monitored weekly and adjusted to pH 7.4 if necessary. After 12 weeks, the AGE-BSA solution is dialyzed against PBS for 48 hours with 4 buffer changes. Protein concentration is determined by the micro-Lowry method.

25 Assay protocol. A Biocoat (Collaborative Biomedical Products) plate is blocked with Superblock (300 μ l /well) at 37°C for 1hr and washed with PBS-Tween three times. AGE-BSA was diluted in PBS to a concentration required to obtain maximum cross-linking as determined in a preliminary experiment. One hundred μ l of the AGE-BSA working solution is added to test wells and a similar concentration of BSA is added to blank wells. The first three wells are left empty for the reagent blank. The plate is incubated at 37°C for four hours and washed with PBS-Tween three times. Test compounds are dissolved in PBS and pH adjusted to 7.0 if necessary. One hundred μ l of

- a test compound is added to triplicate wells. To measure maximum crosslinking 100 μ l of PBS is added to three to six wells. The test compounds and PBS are also added to triplicate wells incubated with BSA, to obtain the blank readings. The plate is then incubated overnight at 37°C. The plate is washed with PBS-Tween, 50 mL of
- 5 Rabbit-anti-BSA antibody (1 to 4000 in PBS) is added to each well and the plate is incubated at room temperature for 60 min. After the plate is washed with PBS-Tween, 50mL of horseradish peroxidase-conjugated -goat-anti-rabbit IgG (1 to 4000 in PBS) is added to each well except the first three wells. The plate is incubated at room temperature for 30 min. and washed with PBS-Tween. Two hundred μ l of ABTS
- 10 substrate (prepared from HRP-substrate buffer 10X (Sigma) diluted 1:10 in deionized water and mixed with 50X ABTS reagent (Sigma)) is added to each well and color developed at 37°C for 15 min. Optical density is read at 410 nm with the sample filter set to "1" and the reference filter set to "5" on the Dynatech ELISA plate reader.

- Data Analysis: The average optical density (OD) is calculated for each triplicate
- 15 determination: Corrected OD = (Average of OD AGE-BSA wells - Average of OD BSA wells). Percent breaking by test compounds is expressed as the percent decrease in OD of TTC-AGE-BSA wells incubated with test compounds compared to TTC-AGE-BSA wells incubated with PBS.


- Using the procedure above, it was found that 3 mM 3-[2-(3-phenyl-5-isoxazolyl)-
- 20 2-oxoethyl]-thiazolium bromide caused a 15% reversal of AGE cross-links.

- This invention may be embodied in other forms or carried out in other ways without departing from the spirit or essential characteristics thereof. The present disclosure is therefore to be considered as in all respects illustrative and not restrictive, the scope of the invention being indicated by the appended Claims, and all changes
- 25 which come within the meaning and range of equivalency are intended to be embraced therein.

Various citations to the literature are presented herein, all of which are incorporated herein in their entireties.

What is claimed:

1. A pharmaceutical composition for administration to an animal, comprising a pharmaceutically effective amount of a compound selected from the group consisting of
5 compounds of the formula:



wherein R^1 and R^2 are independently selected from the group consisting of
10 hydrogen and a alkyl group, which can be substituted by a hydroxy group;

Y is a

group of the formula $-CH_2C(=O)R$ wherein R is a heterocyclic group
other than alkylenedioxyaryl containing 4-10 ring members and 1-
3 heteroatoms selected from the group consisting of oxygen,
15 nitrogen and sulfur, the heterocyclic group can be substituted by
one or more substituents selected from the group consisting of
alkyl, oxo, alkoxycarbonylalkyl, aryl, and aralkyl groups; and said
one or more substituents can be substituted by one or more alkyl
or alkoxy groups; or

20 group of the formula $-CH_2C(=O)-NHR'$ wherein R' is a heterocyclic
group other than alkylenedioxyaryl containing 4-10 ring members
and 1-3 heteroatoms selected from the group consisting of oxygen,
nitrogen, and sulfur, the heterocyclic group can be substituted by
one or more alkoxycarbonylalkyl groups; and

25 X is a pharmaceutically acceptable ion; and
a pharmaceutically acceptable carrier therefor.


2. The composition of Claim 1 wherein Y is a group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{R}$ and wherein the alkyl, alkylene and alkoxy groups comprise up to six carbons .
3. The composition of claim 2, wherein the heterocyclic group is axepinyl, azocinyl, hexamethyleneimino, imidazolyl, isoxazolyl, phenylisoxazolyl, morpholinyl, thiomorpholinyl, piperidiny, homopiperidiny, piperazinyl, pyridyl, pyrimidinyl, tetrahydroquinolyl, pyrrolidinyl, thiazolyl, thiazolinyl, thienyl, thionaphthenyl thiophenyl, or 1,2,4-triazoyl.
4. The composition of claim 3 wherein Y is selected from the group consisting of 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]; 3-(2-(4-morpholinyl)-2-oxoethyl); 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]; 3-(2-(1-piperidiny)-2-oxoethyl); 3-(2-(2-furanyl)-2-oxoethyl); 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinoliny)-2-oxoethyl]; 3-(2-(1-pyrrolidinyl)-2-oxoethyl); 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]; 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]; 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(4-thiomorpholinyl)-2-oxoethyl); 3-(2-(hexahydro-1-azepinyl)-2-oxoethyl); 3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]; 3-(2-(octahydro-1-azocinyl)-2-oxoethyl); 3-(2-(2-pyridinyl)-2-oxoethyl); 3-[2-(2-methyl-1-piperidiny)-2-oxoethyl]; 3-[2-(2,6-dimethyl-1-piperidiny)-2-oxoethyl]; 3-[2-(4-benzyl-1-piperidiny)-2-oxoethyl]; and 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl].
5. The composition of Claim 4 wherein said compound is a 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-thiazolium , 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4,5-dimethylthiazolium , 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4-methyl-5-(2-hydroxyethyl)-thiazolium, 3-(2-(4-morpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium , 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(1-piperidiny)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(2-furanyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide, 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinoliny)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium, 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(2-thienyl)-2-oxoethyl)-4,5-dimethyl-thiazolium, 3-(2-(2-thienyl)-2-oxoethyl)-4-methyl-5-hydroxyethylthiazolium, 3-(2-(4-thiomorpholinyl)-2-

oxoethyl)-4,5-dimethylthiazolium, 3-(2-(hexahydro-1-azepinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(octahydro-1-azocinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(2-pyridinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(2,6-dimethyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(4-benzyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium, or 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium salt.

6. The composition of Claim 1 wherein said compound has the formula wherein Y is group of the formula $-\text{CH}_2\text{C}(=\text{O})-\text{NHR}'$ and wherein the alkyl, alkylene and alkoxy groups comprise up to six carbons.

7. The composition of Claim 6 wherein said compound is a 3-[2-[4-(2-ethoxy-2-oxoethyl)-2-thiazolyl]amino-2-oxoethyl]-4,5-dimethylthiazolium salt.

8. A method of treating diabetes or adverse sequelae of diabetes in an animal, comprising administering an effective amount of composition comprising a compound selected from the group consisting of compounds of the formula



wherein R^1 and R^2 are independently selected from the group consisting of hydrogen and an alkyl group, which can be substituted by a hydroxy group;

Y is a

group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{R}$ wherein R is a heterocyclic group other than alkylenedioxyaryl containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, the heterocyclic group can be substituted by one or more substituents selected from the group consisting of alkyl, oxo, alkoxycarbonylalkyl, aryl, and aralkyl groups; and said one or more substituents can be substituted by one or more alkyl or alkoxy groups; or

group of the formula $-\text{CH}_2\text{C}(=\text{O})-\text{NHR}'$ wherein R' is a heterocyclic group other than alkylenedioxyaryl containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur, the heterocyclic group can be substituted by one or more alkoxycarbonylalkyl groups; and

X is a pharmaceutically acceptable ion; and
a carrier therefor.

9. The method of Claim 8 comprising administering the composition wherein Y is a group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{R}$ and wherein the alkyl, alkylene and alkoxy groups comprise up to six carbons.

10. The method of claim 8, comprising administering the composition wherein the heterocyclic group is axepinyl, azocinyl, hexamethyleneimino, imidazolyl, isoxazolyl, phenylisoxazolyl, morpholinyl, thiomorpholinyl, piperidinyl, homopiperidinyl, piperazinyl, pyridyl, pyrimidinyl, tetrahydroquinolyl, pyrrolidinyl, thiazolyl, thiazolinyl, thienyl, thionaphenyl thiophenyl, or 1,2,4-triazolyl.

11. The method of claim 10 comprising administering the composition wherein Y is selected from the group consisting of 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]; 3-(2-(4-morpholinyl)-2-oxoethyl); 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]; 3-(2-(1-piperidinyl)-2-oxoethyl); 3-(2-(2-furanyl)-2-oxoethyl); 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinolinyl)-2-oxoethyl]; 3-(2-(1-pyrrolidinyl)-2-oxoethyl); 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]; 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]; 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(4-thiomorpholinyl)-2-oxoethyl); 3-(2-

(hexahydro-1-azepinyl)-2-oxoethyl); 3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]; 3-(2-(octahydro-1-azocinyl)-2-oxoethyl); 3-(2-(2-pyridinyl)-2-oxoethyl); 3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]; 3-[2-(2,6-dimethyl-1-piperidinyl)-2-oxoethyl]; 3-[2-(4-benzyl-1-piperidinyl)-2-oxoethyl]; and 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl].


5

12. The method of Claim 11 comprising administering the composition wherein said compound is a 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-thiazolium, 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4-methyl-5-(2-hydroxyethyl)-thiazolium, 3-(2-(4-morpholinyl)-2-oxoethyl)-4,5-
- 10 dimethylthiazolium, 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(1-piperidinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(2-furanyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide, 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinolinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(4-phenyl-1-
- 15 piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(2-thienyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(2-thienyl)-2-oxoethyl)-4-methyl-5-hydroxyethylthiazolium, 3-(2-(4-thiomorpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(hexahydro-1-azepinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(octahydro-1-azocinyl)-2-oxoethyl)-4,5-dimethylthiazolium,
- 20 3-(2-(2-pyridinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(2,6-dimethyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(4-benzyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium, or 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium .

- 25 13. The method of Claim 8 comprising administering the composition wherein said compound has the formula $\text{-CH}_2\text{C(=O)-NHR'}$ and wherein the alkyl, alkylene and alkoxy groups comprise up to six carbons.

14. The method of Claim 13 comprising administering the composition wherein said
- 30 compound is 3-[2-[4-(2-ethoxy-2-oxoethyl)-2-thiazolyl]amino-2-oxoethyl]-4,5-dimethylthiazolium chloride or another biologically acceptable salt thereof.

15. A method of, in a mammal, treating kidney damage, treating damage to blood vasculature, improving the elasticity or reducing wrinkles of the skin, or treating damage to an ocular lens comprising administering an effective amount of a pharmaceutical composition, said pharmaceutical composition comprising a compound selected from the group consisting of compounds of the formula



- wherein R¹ and R² are independently selected from the group consisting of hydrogen and an alkyl group, which can be substituted by a hydroxy group;
- Y is a group of the formula -CH₂C(=O)R wherein R is a heterocyclic group other than alkylenedioxyaryl containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, the heterocyclic group can be substituted by one or more substituents selected from the group consisting of alkyl, oxo, alkoxycarbonylalkyl, aryl, and aralkyl groups; and said one or more substituents can be substituted by one or more alkyl or alkoxy groups; or
- group of the formula -CH₂C(=O)-NHR' wherein R' is a heterocyclic group other than alkylenedioxyaryl containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur, the heterocyclic group can be substituted by one or more alkoxycarbonylalkyl groups; and
- X is a pharmaceutically acceptable ion; and

a carrier therefor.


16. The method of Claim 15 comprising administering the composition wherein Y is a group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{R}$ and wherein the alkyl, alkylene and alkoxy groups
5 comprise up to six carbons.
17. The method of claim 15, comprising administering the composition wherein the heterocyclic group is axepinyl, azocinyl, hexamethyleneimino, imidazolyl, isoxazolyl, phenylisoxazolyl, morpholinyl, thiomorpholinyl, piperidinyl, homopiperidinyl,
10 piperazinyl, pyridyl, pyrimidinyl, tetrahydroquinolyl, pyrrolidinyl, thiazolyl, thiazolinyl, thienyl, thionaphthenyl thiophenyl, or 1,2,4-triazolyl.
18. The method of claim 48 comprising administering the composition wherein Y is selected from the group consisting of 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]; 3-(2-(4-
15 morpholinyl)-2-oxoethyl); 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]; 3-(2-(1-piperidinyl)-2-oxoethyl); 3-(2-(2-furanyl)-2-oxoethyl); 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinoliny)-2-oxoethyl]; 3-(2-(1-pyrrolidinyl)-2-oxoethyl); 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]; 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]; 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(4-thiomorpholinyl)-2-oxoethyl); 3-(2-
20 (hexahydro-1-azepinyl)-2-oxoethyl); 3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]; 3-(2-(octahydro-1-azocinyl)-2-oxoethyl); 3-(2-(2-pyridinyl)-2-oxoethyl); 3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]; 3-[2-(2,6-dimethyl-1-piperidinyl)-2-oxoethyl]; 3-[2-(4-benzyl-1-piperidinyl)-2-oxoethyl]; and 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl].
- 25 19. The method of Claim 18 comprising administering the composition wherein said compound is a 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-thiazolium, 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4-methyl-5-(2-hydroxyethyl)-thiazolium, 3-(2-(4-morpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]-4,5-dimethylthiazolium,
30 3-(2-(1-piperidinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(2-furanyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide, 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinoliny)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium, 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(4-phenyl-1-

- piperaziny]l-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(2-thienyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(2-thienyl)-2-oxoethyl)-4-methyl-5-hydroxyethylthiazolium, 3-(2-(4-thiomorpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(hexahydro-1-azepiny]l)-2-oxoethyl)-4,5-dimethylthiazolium, 3-[2-(4-[2-methoxyphenyl]-1-piperaziny]l)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(octahydro-1-azociny]l)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(2-pyridiny]l)-2-oxoethyl)-4,5-dimethylthiazolium, 3-[2-(2-methyl-1-piperidiny]l)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(2,6-dimethyl-1-piperidiny]l)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(4-benzyl-1-piperidiny]l)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(4-benzyl-1-piperaziny]l)-2-oxoethyl]-4,5-dimethylthiazolium, or another biologically acceptable salt thereof.

20. The method of Claim 17 comprising administering the composition wherein said compound has the formula wherein Y is a group of the formula $-\text{CH}_2\text{C}(=\text{O})-\text{NHR}'$ and wherein the alkyl, alkylene and alkoxy groups comprise up to six carbons.

21. The method of Claim 20 comprising administering the composition wherein said compound is a 3-[2-[4-(2-ethoxy-2-oxoethyl)-2-thiazolyl]amino-2-oxoethyl]-4,5-dimethylthiazolium salt.

22. A compound of the formula



wherein R^1 and R^2 are independently selected from the group consisting of hydrogen and an alkyl group, which can be substituted by a hydroxy group;

Y is a

group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{R}$ wherein R is a heterocyclic group other than alkylenedioxyaryl containing 4-10 ring members and one heteroatom selected from the group consisting of nitrogen and sulfur, or 2-3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, the heterocyclic group can be substituted by one or more substituents selected from the group consisting of alkyl, oxo, alkoxycarbonylalkyl, aryl, and aralkyl groups; and said one or more substituents can be substituted by one or more alkyl or alkoxy groups, with the proviso that R is not furanyl; or

group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{NHR}'$ wherein R' is a heterocyclic group other than alkylenedioxyaryl containing 4-10 ring members and 1-3 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur, the heterocyclic group can be substituted by one or more alkoxycarbonylalkyl groups;

and X is a pharmaceutically acceptable ion.

23. The compound of claim 22 wherein said compound has the formula wherein Y is group of the formula $-\text{CH}_2\text{C}(=\text{O})\text{R}$ and wherein the alkyl, alkylene and alkoxy groups comprise up to six carbons.

24. The compound of claim 22, wherein the heterocyclic group is axepinyl, azocinyl, hexamethyleneimino, imidazolyl, isoxazolyl, phenylisoxazolyl, morpholinyl, thiomorpholinyl, piperidinyl, homopiperidinyl, piperazinyl, pyridyl, pyrimidinyl, tetrahydroquinolyl, pyrrolidinyl, thiazolyl, thiazolinyl, thienyl, thionaphenyl thiophenyl, or 1,2,4-triazolyl.

25. The compound of claim 24 wherein Y is selected from the group consisting of 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]; 3-(2-(4-morpholinyl)-2-oxoethyl); 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]; 3-(2-(1-piperidinyl)-2-oxoethyl); 3-(2-(2-furanyl)-2-oxoethyl); 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinolinyl)-2-oxoethyl]; 3-(2-(1-pyrrolidinyl)-2-

oxoethyl; 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]; 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]; 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(2-thienyl)-2-oxoethyl); 3-(2-(4-thiomorpholinyl)-2-oxoethyl); 3-(2-(hexahydro-1-azepinyl)-2-oxoethyl); 3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]; 3-(2-(octahydro-1-azocinyl)-2-oxoethyl); 3-(2-(2-pyridinyl)-2-oxoethyl); 3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]; 3-[2-(2,6-dimethyl-1-piperidinyl)-2-oxoethyl]; 3-[2-(4-benzyl-1-piperidinyl)-2-oxoethyl]; and 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl].

26. The compound of Claim 25 wherein said compound is a 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-thiazolium, 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(3-phenyl-5-isoxazolyl)-2-oxoethyl]-4-methyl-5-(2-hydroxyethyl)-thiazolium, 3-(2-(4-morpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-[2-(2,6-dimethyl-4-morpholinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(1-piperidinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(2-furanyl)-2-oxoethyl)-4,5-dimethylthiazolium bromide, 3-[2-(2-oxo-1,2,3,4-tetrahydro-6-quinolinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(1-pyrrolidinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium, 3-[2-(3-methyl-2-thianaphthenyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(4-phenyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(2-thienyl)-2-oxoethyl)-4,5-dimethyl-thiazolium, 3-(2-(2-thienyl)-2-oxoethyl)-4-methyl-5-hydroxyethylthiazolium, 3-(2-(4-thiomorpholinyl)-2-oxoethyl)-4,5-dimethylthiazolium, 3-(2-(hexahydro-1-azepinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium, 3-[2-(4-[2-methoxyphenyl]-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-(2-(octahydro-1-azocinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium, 3-(2-(2-pyridinyl)-2-oxoethyl)-4,5-dimethyl-thiazolium, 3-[2-(2-methyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(2,6-dimethyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium, 3-[2-(4-benzyl-1-piperidinyl)-2-oxoethyl]-4,5-dimethylthiazolium, or 3-[2-(4-benzyl-1-piperazinyl)-2-oxoethyl]-4,5-dimethylthiazolium.

27. The compound of Claim 22 wherein said compound has the formula wherein Y is group of the formula $-\text{CH}_2\text{C}(=\text{O})-\text{NHR}'$ and wherein the alkyl, alkylene and alkoxy groups comprise up to six carbons.

28. The compound of Claim 27 wherein said compound is a 3-[2-(2'-amino-5'-carboethoxymethylene-thiazolyl)-2-oxoethyl]-4,5-dimethyl-thiazolium salt.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/26565

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/427; C07D 417/06

US CL : 548/204; 514/365

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 548/204; 514/365

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TSUGE O. Stereochemical study on 1,3 dipolar cycloaddition reactions of heteroaromatic n-ylides with symmetrically substituted cis and trans olefins bull.Chem. Soc. Japan, 1985 vol.58, pages 3137-3158 1985 SEE CPD 6b.	1,2,22,23

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

11 JANUARY 2000

Date of mailing of the international search report

14 MAR 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ROBERT GERSTL

Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet)(July 1992)*

